



GROUP 7 3 4 FUNGICIDES

EverGol Energy

Seed Treatment Fungicide

A Seed Treatment Fungicide for seed and seedling protection against certain seed-borne and soil-borne diseases on labelled crops.

All seed treated with this product must be conspicuously coloured at the time of treatment. No open transfer is permitted in commercial seed treatment facilities.

SUSPENSION

AGRICULTURAL

GUARANTEE:	Penflufen	38.4 g/L
	Prothioconazole	76.8 g/L
	Metalaxyl	61.4 g/L

Contains 1,2 benzisothiazolin-3-one at 0.05% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 30364 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

PROTECT FROM FREEZING

NET CONTENTS: 0.946L - 1000L

Bayer CropScience Inc.
Suite 200, 160 quarry Park Blvd. SE
Calgary, AB T2C 3G3

For product information, call: 1-888-283-6847
In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

EverGol Energy is a seed treatment containing the fungicides penflufen, prothioconazole and metalaxyl. EverGol Energy provides protection from seed, seedling and soil-borne diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Pythium* spp., *Botrytis cinerea* and *Phomopsis longicolla* on labelled crops.

DIRECTIONS FOR USE:

Refer to the attached booklet for specific use directions for each crop and detailed instructions on rates, diseases controlled, application timing and technique.

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Commercial seed treatment facilities must use closed transfer equipment only including closed mixing, loading, calibrating, and treatment equipment. No open transfer of EverGol Energy is permitted in commercial seed treatment facilities.
- Workers must wear long pants, a long-sleeved shirt and chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, bagging, sewing or stacking of bagged treated seed, or handling treated seed. In addition, workers must wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.
- Do not apply in a way that this product will contact workers or other persons. Only handlers (mixers, loaders and applicators) wearing personal protective equipment may be in the area being treated during application.
- Use good personal hygiene, washing hands and exposed skin with warm water and soap before eating, drinking, or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on the acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. Medical Personnel should contact Bayer's Medical Information Services, Toll-Free: 1-800-334-7577.

ENVIRONMENTAL HAZARDS:

Toxic to aquatic organisms and non-target terrestrial plants. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other waters.

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. DO NOT leave exposed treated seed on soil surface. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water is shallow.

STORAGE:

To prevent contamination store this product away from food or feed. Store in a cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 40°C or go below -10°C.

DISPOSAL:

FOR NON-RETURNABLE CONTAINERS:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

1. Transfer EverGol Energy from the container to the seed treating equipment applicator tank.
2. Add 1/3 of the rinse water to the container and agitate the contents.
3. Transfer the rinsate to the applicator tank.
4. Rinse as above 2 more times.
5. Thoroughly mix the EverGol Energy and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.



EverGol Energy

Seed Treatment Fungicide

GROUP	7 3 4	FUNGICIDES
-------	-------	------------

A Seed Treatment Fungicide for seed and seedling protection against certain seed-borne and soil-borne diseases on labelled crops.

All seed treated with this product must be conspicuously coloured at the time of treatment. No open transfer is permitted in commercial seed treatment facilities.

SUSPENSION

AGRICULTURAL

GUARANTEE:	Penflufen	38.4 g/L
	Prothioconazole	76.8 g/L
	Metalaxyl	61.4 g/L

Contains 1,2 benzisothiazolin-3-one at 0.05% and 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0009% and 2-methyl-4-isothiazolin-3-one at 0.0003% as preservatives.

REGISTRATION NO. 30364 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

PROTECT FROM FREEZING

NET CONTENTS: 0.946L - 1000L

Bayer CropScience Inc.
Suite 200, 160 quarry Park Blvd. SE
Calgary, AB T2C 3G3

For product information, call: 1-888-283-6847

In case of spills, poisoning, fire or other emergencies,
call: 1-800-334-7577 (24 hours a day)

Table of Contents:	Section Number
GENERAL INFORMATION	The Product 1
SAFETY AND HANDLING	Precautions, Protective Clothing and Equipment 2
	First Aid and Toxicological Information 3
	Environmental Hazards 4
	Storage 5
	Disposal 6
	Notices 7
DIRECTIONS FOR USE	Pests Controlled 8
	Recommended Uses 9
	Application Cautions 10
	Use Limitations 11
	Resistance Management Recommendations 12

GENERAL INFORMATION

Section 1: The Product

EverGol Energy is a seed treatment containing the fungicides penflufen, prothioconazole and metalaxyl. EverGol Energy provides protection from seed, seedling and soil-borne diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Pythium* spp., *Botrytis cinerea* and *Phomopsis longicolla* on labelled crops.

SAFETY AND HANDLING

Section 2: Precautions, Protective Clothing and Equipment

PRECAUTIONS:

- KEEP OUT OF REACH OF CHILDREN.
- Commercial seed treatment facilities must use closed transfer equipment only including closed mixing, loading, calibrating, and treatment equipment. No open transfer of EverGol Energy is permitted in commercial seed treatment facilities.
- Workers must wear long pants, a long-sleeved shirt and chemical-resistant gloves during mixing, loading, treating, clean-up, maintenance of seed treatment equipment, bagging, sewing or stacking of bagged treated seed, or handling treated seed. In addition, workers must wear a suitable dust mask when bagging or sewing bags of treated seed or when transferring seed to a storage bin.
- Do not apply in a way that this product will contact workers or other persons. Only handlers (mixers, loaders and applicators) wearing personal protective equipment may be in the area being treated during application.
- Use good personal hygiene, washing hands and exposed skin with warm water and soap before eating, drinking, or smoking. No food, drink or tobacco should be allowed in areas of chemical storage or use.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on the acceptable residue levels in the U.S., visit CropLife Canada's website at www.croplife.ca.

Section 3: First Aid and Toxicological Information

FIRST AID:

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove

contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically. Medical Personnel should contact Bayer's Medical Information Services, Toll-Free: 1-800-334-7577.

Section 4: Environmental Hazards

ENVIRONMENTAL HAZARDS:

Toxic to aquatic organisms and non-target terrestrial plants. DO NOT discharge effluent containing this product into sewer systems, lakes, streams, ponds, estuaries, oceans or other waters.

Dispose of all excess treated seed. Left over treated seed may be double-sown around the headland or buried away from water sources in accordance with local requirements. DO NOT leave exposed treated seed on soil surface. Any spilled or exposed seeds must be incorporated into the soil or otherwise cleaned-up from the soil surface.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water is shallow.

Section 5: Storage

To prevent contamination store this product away from food or feed. Store in a cool, dry area. Do not store in direct sunlight. Do not allow prolonged storage in temperatures that exceed 40°C or go below -10°C.

Section 6: Disposal

FOR NON-RETURNABLE CONTAINERS:

5. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.

6. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
7. Make the empty container unsuitable for further use.
8. Dispose of the container in accordance with provincial requirements.

FOR RECYCLABLE CONTAINERS:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

3. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
4. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

FOR RETURNABLE CONTAINERS:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

FOR REFILLABLE CONTAINERS:

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

TRIPLE RINSE INSTRUCTIONS (FOR RECYCLABLE AND DISPOSABLE CONTAINERS ONLY): This container should be triple-rinsed prior to its disposal. The rinsate can be added to the seed treatment. Follow these steps for rinsing of containers:

6. Transfer EverGol Energy from the container to the seed treating equipment applicator tank.
7. Add 1/3 of the rinse water to the container and agitate the contents.
8. Transfer the rinsate to the applicator tank.
9. Rinse as above 2 more times.
10. Thoroughly mix the EverGol Energy and rinse water in the applicator tank prior to use.

CAUTION: Be sure to adjust the undiluted application rate by the same volume of water added to compensate for the dilution of the product caused by the addition of rinsate.

DISPOSAL OF UNUSED, UNWANTED PRODUCT:

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

Section 7: Notices

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

DIRECTIONS FOR USE

DO NOT apply this product or treated seeds directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches and wetlands), estuaries or marine habitats.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Section 8: Pests Controlled

EverGol Energy is a seed treatment containing the fungicides penflufen, prothioconazole and metalaxyl. EverGol Energy provides seed and seedling protection against a wide range of diseases caused by *Rhizoctonia solani*, *Fusarium* spp., *Pythium* spp., *Botrytis cinerea* and *Phomopsis longicolla* on labelled crops.

Section 9: Recommended Uses

EverGol Energy is a seed treatment formulation for use in commercial seed treatment operations, and for on-farm treating with conventional seed treating equipment which can accurately meter and apply flowable seed treatment formulations.

This product is recommended to be diluted with water or another suitable liquid just prior to application to ensure uniform coverage on the seed during the application process. Uniform application to seed is necessary to ensure optimum performance. Allow seeds to dry before bagging, storing or seeding.

CROP	PEST	APPLICATION RATE	DIRECTIONS
Wheat, barley, oat, rye, triticale, pearl millet, proso millet	<p>Diseases Controlled:</p> <p>Loose smut (<i>Ustilago tritici</i>)</p> <p>Common bunt (<i>Tilletia caries</i>, <i>Tilletia laevis</i>)</p> <p>True Loose smut (<i>Ustilago nuda</i>)</p> <p>False loose smut (<i>Ustilago nigra</i>)</p> <p>Covered smut (<i>Ustilago hordei</i>)</p> <p>Leaf stripe (<i>Pyrenophora graminea</i>)</p> <p>Loose smut (<i>Ustilago avenae</i>)</p> <p>Covered smut (<i>Ustilago. kolleri</i>)</p> <p>Seed rot / pre-emergence damping-off and seedling blight caused by seed-borne <i>Fusarium</i> spp. and <i>Cochliobolus sativus</i></p> <p>Seed rot / pre-emergence damping-off caused by soil-borne <i>Fusarium</i> spp., <i>Pythium</i> spp. and <i>Cochliobolus sativus</i></p> <p>Post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp.</p> <p>Seed rot / pre-emergence damping-off, post-emergence damping-off and seedling blight caused by seed-borne <i>Aspergillus</i> spp.</p>	65 mL per 100 kg of seed	<p>EverGol Energy is designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide a good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure optimum product performance.</p> <p>This product contains no dye. An appropriate seed colourant must be applied.</p>
	<p>Diseases Suppressed:</p> <p>Root rot caused by seed-borne and soil-borne <i>Fusarium</i> spp.</p> <p>Root rot caused by soil-borne <i>Cochliobolus sativus</i></p> <p>Crown rot caused by seed-borne <i>Fusarium</i> spp.</p> <p>Seedling blight caused by seed-borne <i>Penicillium</i> spp.</p>		

CROP	PEST	APPLICATION RATE	DIRECTIONS
Corn (field, sweet, popcorn)	<p>Diseases Controlled:</p> <p>Seed rot / pre-emergence damping-off caused by seed-borne and soil-borne <i>Fusarium</i> spp.</p> <p>Post-emergence damping-off caused by soil-borne <i>Fusarium</i> spp.</p> <p>Seed rot / pre-emergence damping off caused by soil-borne <i>Pythium</i> spp.</p> <p>Seed rot / pre-emergence damping off caused by <i>Rhizoctonia solani</i></p> <p>Seed rot / pre-emergence damping-off caused by seed-borne <i>Cladosporium</i> spp. and <i>Aspergillus</i> spp.</p>	65 mL per 100 kg of seed	<p>EverGol Energy is designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide a good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure optimum product performance.</p> <p>This product contains no dye. An appropriate seed colourant must be applied.</p>
	<p>Diseases Suppressed:</p> <p>Seed rot / pre-emergence damping-off caused by <i>Penicillium</i> spp.</p>		

CROP	PEST	APPLICATION RATE	DIRECTIONS
<p>Crop SubGroup 6C (dry shelled pea and beans), and soybean:</p> <p>Soybean, lupin, sweet lupin, white lupin and white sweet lupin, field bean, kidney bean, lima bean (dry), navy bean, pinto bean, tepary bean, adzuki bean, blackeyed pea, catjang, cowpea, Crowder pea, moth bean, mung bean, rice bean, southern pea, curd bean, broad bean, chickpea, guar, lablab bean, lentil, field pea, pigeon pea</p>	<p>Diseases Controlled:</p> <p>Seed rot / pre-emergence damping-off and post-emergence damping-off caused by <i>Rhizoctonia solani</i>, <i>Fusarium</i> spp., and <i>Pythium</i> spp.</p> <p>Early-season root rot and seedling blight caused by <i>Rhizoctonia solani</i> and <i>Fusarium</i> spp. Seedling blight caused by seed-borne <i>Botrytis cinerea</i></p> <p>Seed rot / pre-emergence damping-off of soybean caused by <i>Phomopsis longicolla</i></p> <hr/> <p>Diseases Suppressed:</p> <p>Seed rot / pre-emergence damping-off of chickpea caused by seed-borne <i>Ascochyta rabiei</i></p>	<p>65 mL per 100 kg of seed</p>	<p>EverGol Energy is designed for commercial or on-farm treating with conventional seed treating equipment which can accurately control application rates and provide a good distribution of the chemical onto the seed in the mixing chamber.</p> <p>Uniform application to seed is necessary to ensure optimum product performance.</p> <p>This product contains no dye. An appropriate seed colourant must be applied.</p>

TANK-MIXTURES

For control of certain insect pests and/or additional diseases not on the EverGol Energy product label, EverGol Energy may be combined with certain seed treatment insecticides and fungicides.

Refer to the registered label of the tank-mix partner(s) for application rates, precautions and directions for use associated with those products, and follow the most restrictive label precautions and limitations.

CROP	TANK-MIX PARTNER
Soybean ¹	Stress Shield for cereals and soybeans
Corn (field, sweet, popcorn)	Poncho 600 FS
	Trilex FS
Wheat, barley, oat	Stress Shield for cereals and soybeans
	Stress Shield for cereals

¹ For control of early-season *Phytophthora* in soybean, EverGol Energy may also be tankmixed with 35 mL of Allegiance FL per 100 kg seed.

Section 10: Application Cautions

COLOURATION OF TREATED SEED:

This product contains no colourant. An appropriate colourant must be added when this product is applied to seed. Regulations pertaining to colouration of treated seed enforced under the *Seeds Act* must be strictly adhered to when using this product.

SLURRY PREPARATION:

When preparing a slurry of EverGol Energy with dilutants such as water or liquid inoculants, additional or continuous agitation or mixing may be necessary to prevent EverGol Energy from settling out. Do not store diluted slurry mixtures.

SEED TREATMENT AND INOCULANTS:

Please check with inoculant manufacturers for application and compatibility details prior to use.

Seeds treated with EverGol Energy and seed inoculants may not flow through seeding equipment at the same rate as untreated seeds. Recalibrate the seeding equipment prior to planting treated seeds. Mixing EverGol Energy with seed inoculants may increase drying time while treating and extend the processing time.

SEED QUALITY:

Lab and field studies have shown that EverGol Energy treated seed can be stored without loss in

germination or disease performance. Many large-seed legumes are prone to mechanical damage during seed handling, and a reduction in germination within a seed lot can occur should handling cause seed damage irrespective of seed treatment application. EverGol Energy-treated or untreated seed may drop in germination with age and germination drops are accelerated under adverse storage conditions or in seed with a high level of mechanical damage. Use handling equipment designed to minimize mechanical damage when moving seed. Due to seed quality and seed storage conditions beyond the control of Bayer CropScience, no claims are made to guarantee the germination of carry-over seed or propagating materials for all crop seed. Treatment of highly mechanically scarred or damaged seed, or seed known to be of low vigour and poor quality, may result in reduced germination and/or reduction of seed and seedling vigour. Treat a small quantity of seed using equipment similar to that planned for treating the total seed lot. Conduct germination tests on a small portion of seed before committing the total seed lot to a selected seed treatment.

Section 11: Use Limitations

USE RESTRICTIONS:

- Treated seed must not be used for food, feed or oil processing.
- Do not apply this product directly to freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, ditches, and wetlands), estuaries, or marine habitats.
- Do not contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

LABELLING TREATED SEED:

All bags containing treated seed must be labelled or tagged as follows: “This seed has been treated with EverGol Energy, which contains penflufen, prothioconazole and metalaxyl. When handling treated seed wear a long-sleeved shirt, long pants and chemical-resistant gloves. DO NOT use for feed, food or oil processing. Store away from feeds and other foodstuffs.”

ROTATIONAL CROPS:

Registered crops for EverGol Energy, as well as canola, mustard, rapeseed, borage, flax, crambe and potato, may be replanted at any time. For all other crops, do not plant-back within 30 days of seeding with EverGol Energy-treated seed.

Section 12: Resistance Management Recommendations

For resistance management, please note that EverGol Energy contains a Group 3, 4, and 7 fungicide. Any fungal population may contain individuals naturally resistant to EverGol Energy and other Group 3, 4 and 7 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may

also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of EverGol Energy or other Group 3, 4 and 7 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group when such use is permitted.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
 - For further information or to report suspected resistance contact Bayer CropScience Inc at 1-888-283-6847 or at www.bayercropscience.ca.