



# THE BROADLEAF WEED CONTROL YOU NEED WITH EXCEPTIONAL CROP SAFETY

# LAUDIS<sup>®</sup>

## REGISTERED TANK MIXES:

- Roundup Transorb<sup>®</sup> HC liquid herbicide<sup>1</sup>
- Roundup WeatherMAX<sup>®</sup> with Transorb<sup>®</sup> 2 Technology liquid herbicide<sup>1</sup>
- Roundup Xtend<sup>®</sup> herbicide with VaporGrip<sup>®</sup> Technology<sup>1,2</sup>
- XtendiMax<sup>®</sup> herbicide with VaporGrip<sup>®</sup> Technology<sup>2</sup>
- Atrazine
- Pardner<sup>®</sup> herbicide

## THE PRODUCT

Laudis<sup>®</sup> herbicide provides fast acting broadleaf weed control (including glyphosate resistant biotypes) with short-term residual activity and excellent crop safety in field corn and sweet corn.

## THE LAUDIS ADVANTAGE

- Excellent post emergence weed control, including tough to control glyphosate-resistant weeds, such as kochia, giant ragweed and waterhemp
- Built-in safener for exceptional crop safety on field corn and sweet corn
- Rapid burndown and up to 3 weeks of residual activity at the high rate 220 mL/ha (89 mL/ac.)
- Excellent resistance management tool and tank mix partner with Roundup<sup>®</sup> brands
- Consistent and fast weed control under a variety of conditions


<sup>1</sup> Use in corn hybrids with Roundup Ready<sup>®</sup> 2 Technology only.

<sup>2</sup> Do not tank mix these products with UAN or an additional adjuvant.

<b>ACTIVE INGREDIENT</b>	Tembotrione (Group 27) Crop safener - isoxadifen
<b>FORMULATION</b>	Suspension concentrate
<b>CROPS FOR USE</b>	Corn (field and sweet)
<b>RAINFASTNESS</b>	Rainfast 2 hours after application
<b>WATER VOLUME</b>	Apply in a minimum of 100 L/ha (10 gal./ac.). For weed control in dense weed populations or under adverse growing conditions, 150 to 200 L/ha (15 to 20 gal./ac.) is recommended.
<b>PACKAGING</b>	3.6 L jug
<b>RATE</b>	145 mL/ha (59 mL/ac.) (low rate) – (60 ac./jug) 220 mL/ha (89 mL/ac.) (high rate)* – (40 ac./jug)


\*refer to detailed application rate table for more information

## LAUDIS IN A 2-PASS PROGRAM



Source: 2021 Western Canada Market Development (n=5). A total of 17 weed hits, 13 broadleaves and 4 grasses. Treatment means are significantly different at P<0.05. Your results may vary depending on agronomic, environmental and pest pressure variables.

## LAUDIS IN A POST PROGRAM



Source: 2021 Western Canada Market Development (n=8). A total of 21 broadleaves and 6 grasses. Treatment means are significantly different at P<0.05. Your results may vary depending on agronomic, environmental and pest pressure variables.

WEEDS CONTROLLED <sup>1</sup>	RECOMMENDED STAGE	PRODUCTS	RATES
Common lamb's-quarters	Up to 6-leaf stage	Laudis® herbicide plus adjuvant system*	145 mL/ha (59 mL/ac.)
Giant foxtail <sup>2</sup>	Up to 2 tillers		
Redroot pigweed	Up to 6-leaf stage		
Velvetleaf			
Canada fleabane <sup>3</sup>	Up to 10 cm height/diameter		220 mL/ha (89 mL/ac.)
Common ragweed	Up to 6-leaf stage		
Common waterhemp			
Kochia			
Giant ragweed			
Green foxtail <sup>2</sup>	Up to 2 tillers		
Wild buckwheat <sup>2,4</sup>	Up to 6-leaf stage		

### CROP STAGING: SEQUENTIAL APPLICATION OPTION

Crop Stage at Application Timing	Field corn*	Sweet corn*
1	2- to 5-leaf stage	2-leaf stage up to and including the 8-leaf stage
2	up to and including the 8-leaf stage	Do not apply

### RE-CROP INTERVAL

Immediate Plant Back	3-4 months	10 months	22 months
Field corn Sweet corn	Winter wheat	Potatoes Soybeans Spring wheat	Dry beans

<sup>1</sup> Includes ALS/SU (Group 2), Auxin (Group 4), Glyphosate (Group 9), PPO (Group 14) resistant biotypes, and Triazine (Group 5).

<sup>2</sup> Suppression only

<sup>3</sup> For improved control of Canada fleabane - field corn only: apply Laudis herbicide in tank-mix with one of the dicamba-containing tank-mix partners. Refer to label for more details.

<sup>4</sup> For control of wild buckwheat - field corn only: Laudis herbicide may be tank-mixed with 1.2 L/ha of Aatrex® Liquid 480 Herbicide. Maximum of one (1) application per season. Apply to field corn at the 2- to 5-leaf stage.

\* Do not apply more than two applications of Laudis herbicide to field corn or more than one application to sweet corn, per growing season

### APPLICATION TIPS

- Laudis is to be used in conjunction with a Methylated Seed Oil (MSO) based surfactant
- For dicamba-containing tank-mix partners, follow label recommendations
- Use of a spray-grade liquid nitrogen fertilizer is recommended
- Restricted Entry Interval (REI) is 12 hours
- If tank-mixing, always respect the maximum pre-harvest/-grazing interval stated on the labels of all the tank-mix products.
- Pre-harvest interval: 45 days for grazing and harvested grain or forage.



cropscience.bayer.ca | 1 888-283-6847 | @Bayer4CropsCA | #AskBayerCrop

Bayer is a member of Excellence Through Stewardship® (ETS). Bayer products are commercialized in accordance with ETS Product Launch Stewardship Guidance, and in compliance with Bayer's Policy for Commercialization of Biotechnology-Derived Plant Products in Commodity Crops. These products have been approved for import into key export markets with functioning regulatory systems. Any crop or material produced from these products can only be exported to, or used, processed or sold in countries where all necessary regulatory approvals have been granted. It is a violation of national and international law to move material containing biotech traits across boundaries into nations where import is not permitted. Growers should talk to their grain handler or product purchaser to confirm their buying position for these products. Excellence Through Stewardship® is a registered trademark of Excellence Through Stewardship.

**ALWAYS READ AND FOLLOW PESTICIDE LABEL DIRECTIONS.** Roundup Ready® Technology contains genes that confer tolerance to glyphosate. Roundup Ready® 2 Technology contains genes that confer tolerance to glyphosate. Glyphosate will kill crops that are not tolerant to glyphosate. Bayer, Bayer Cross, Laudis®, Pardner®, Roundup®, Roundup Ready 2 Technology and Design™, Roundup Ready®, Roundup Transorb®, Roundup WeatherMAX®, Roundup Xtend®, Transorb®, VaporGrip® and XtendiMax® are trademarks of Bayer Group. Used under license. All other trademarks are the trademarks of their respective companies. Bayer CropScience Inc. is a member of CropLife Canada. ©2022 Bayer Group. All rights reserved.